

On the equivalence between the Barkhausen effect and directed Abelian sandpile models

Alexei Vázquez and Oscar Sotolongo-Costa

Department of Theoretical Physics, Faculty of Physics, Havana University, Havana 10400, Cuba

The existence of self-organized criticality in the Barkhausen effect and its analogy with sandpile models is investigated. It is demonstrated that a model recently introduced to describe the dynamics of a domain wall [Cizeau *et al*, Phys. Rev. Lett. **79**, 4669 (1997)] belongs to the universality class of undirected Abelian sandpile models. In this way it is shown that the Barkhausen effect can be taken as an experimental observation of self-organized critical phenomena.

75.60.Ej, 05.40.+j, 64.60.Lx, 68.35.Rh

One of the fundamental tasks of the theory of critical phenomena is to determine the different universality classes. If different systems or phenomena cannot be grouped into a reduced group of universality classes then the central idea of critical phenomena, the existence of universal behavior, will be of no relevance. This task has been carry out succesfully in ordinary critical phenomena. However, the precise identification of the universality classes in a non-equilibrium critical phenomena like self-organized criticality (SOC) is still unresolved.

SOC was introduced to explain the critical behavior of a vast class of driven dissipative system which evolve into a critical state [1]. In its early state it was believed that such a critical state is insensitive to changes in control parameters and no fine-tuning is needed. More recent interpretations of this phenomena have shown that criticality in SOC systems is obtained after some control parameters, for instance the driving and dissipation rates, are fine-tuned to zero [2].

The Barkhausen effect has been taken as an experimental observation of SOC behavior. Based on phenomenological analogies between the Barkhausen effect and sandpile models, such as the existence of power law distributions of avalanche size and duration, some authors have claim that the Barkhausen effect exhibits SOC behavior [3,4]. However, this conclusion has been criticized by other researches which pointed out that the observation of power law distributions is not necessarily an evidence of SOC behavior [5,6]. There are alternative approaches, like the random field Ising model [7–9], where the power law distributions are a consequence of the scaling properties of disorder. The determination of the universality classes in this case thus become more difficult because it is still not clear if the Barkhausen effect exhibits or not SOC behavior.

Some light in this controversy has been given by Cizeau, Zapperi, Durin and Stanley (CZDE) [10]. They introduced an equation of motion for a single domain wall where dipolar interactions, demagnetization effects and quenched disorder are considered, containing some previous works [11,12] as limiting cases. CZDE observed

that their model has certain analogies with sandpile models. More precisely the critical state is obtained when the magnetic field rate (the driving rate in sandpile models) and the demagnetization factor (the dissipation rate in sandpile models) are fine-tuned to zero. However their analysis, in relation to this analogy, was limited to this phenomenological observation based on numerical simulations and mean field analysis. The great importance of this conjecture is that if one could map the Barkhausen effect into certain class of sandpile models then one could be sure about the existence of SOC in this phenomena. This is precisely the scope of this work.

We investigate the CZDE model in the case of strong magnetization, where dipolar interactions are relevant [13]. First we show that when the magnetization field increases at constant rate the domain wall is never pinned but moves in average at constant velocity. Then we obtain exact expressions for the average interface velocity and susceptibility. The scaling exponents are obtained using perturbative analysis, some of them but not all results identical to those obtained by CZDE for the case of constant magnetic field. Further RG calculations reveals that perturbation theory is exact up to $\mathcal{O}(r^{-1})$, where r is proportional to the square of the saturation magnetization. From the comparison of the scaling exponents with those observed in sandpile models we conclude that the CZDE model with magnetic field increasing at constant rate belongs to the universality class of directed Abelian sandpile models (DASM) [14].

To start our analysis let us introduce the CZDE model. The domain wall is modeled by a d -dimensional interface, dividing two regions of opposite magnetization, moving in a $d + 1$ environment described by its position $h(\vec{x}, t)$. Considering the contribution of magnetostatic, ferromagnetic and magneto-chrystalline interactions one obtains the following equation of motion [10]

$$\lambda \frac{\partial}{\partial t} h(\vec{x}, t) = \Gamma \nabla^2 h(\vec{x}, t) + 2\mu_0 M_s H + \eta[\vec{x}, h(\vec{x}, t)] - 4\mu_0 \mathcal{N} M_s^2 \int \frac{d^d x'}{L^d} h(\vec{x}', t)$$

$$+ \int d^d x' K(\vec{x} - \vec{x}') [h(\vec{x}', t) - h(\vec{x}, t)], \quad (1)$$

where λ is a viscosity coefficient, Γ is the surface tension of the wall, H is the magnetic field intensity, L is the linear size of the system, and M_s is the saturation magnetization per unit volume. Long-range demagnetization effects are described by the fourth term in the right hand side, where $\mathcal{N}$ is the demagnetization factor. Dipolar interactions are characterized by the fifth term, where the kernel $K(\vec{x})$ is anisotropic and has Fourier transform

$$\tilde{K}(\vec{k}) = \frac{\mu_0 M_s^2}{4\pi^2} |\vec{k}| \cos^2 \theta, \quad (2)$$

where θ is the angle between $\vec{k}$ and the magnetization. $\eta(\vec{x}, h)$ is a Gaussian uncorrelated noise due to lattice defects or other factors, with zero mean and noise correlator

$$\langle \eta(\vec{x}, h) \eta(\vec{x}', h') \rangle = \delta^d(\vec{x} - \vec{x}') \Delta(h - h'), \quad (3)$$

where $\Delta(h)$ is a monotonically decreasing even function.

If dipolar interactions and demagnetization effects are neglected and H is constant then eq. (1) is reduced to the Edwards-Wilkinson equation with quenched noise. This limiting case has been extensively studied in the literature [15,16]. A depinning transition takes place at certain critical field H_c , determined by the disorder. For $H < H_c$ the interface is pinned after certain finite time while for $H > H_c$ it moves with finite average velocity. The upper critical dimension is $d_c = 4$. This features remains if dipolar interactions are considered but the upper critical dimension is reduced to $d_c = 2$ [10].

When the demagnetization field is included and the magnetic field increases at rate c then the interface is never pinned by impurities, but always moves with a finite average velocity v . A perturbative solution of eq. (1) can thus be found expanding $h(\vec{x}, t)$ around the flat co-moving interface vt . Taking $h(\vec{x}, t) = vt + y(\vec{x}, t)$, with $\langle y(\vec{x}, t) \rangle = 0$, we obtain the following equation for $y(\vec{x}, t)$

$$\begin{aligned} \lambda \frac{\partial}{\partial t} y(\vec{x}, t) &= \Gamma \nabla^2 y(\vec{x}, t) + 2\mu_0 M_s (c - 2\mathcal{N} M_s v) t \\ -\lambda v + \eta[\vec{x}, vt + y(\vec{x}, t)] &- 4\mu_0 \mathcal{N} M_s^2 \int \frac{d^d x'}{L^d} y(\vec{x}', t) \\ &+ \int d^d x' K(\vec{x} - \vec{x}') [y(\vec{x}', t) - y\vec{x}, t)], \end{aligned} \quad (4)$$

The average velocity is obtained using the constraint $\langle y(\vec{x}, t) \rangle = 0$. For this purpose is better to work with the equation for the Fourier transform of $h(\vec{x}, t)$, $\tilde{h}(\vec{k}, \omega)$. The effective external field $(c - 2\mathcal{N} M_s v)t$ gives a singular term of the order of ω^{-2} . This singular term predominates in the low frequency limit resulting, after imposing $\langle \tilde{y}(\vec{k}, \omega) \rangle = 0$,

$$v = \frac{c}{2\mathcal{N} M_s}. \quad (5)$$

This result is valid to all orders of perturbation expansion and, therefore, exact. In the MF theory by CZDE, which is equivalent to the ABBM model, it is obtained that $v \sim c/\mathcal{N} M_s$ in agreement with eq. (5). We have thus shown that this result is exact and, therefore, valid beyond the MF theory.

Another exact result can be obtained if one computes the low-frequency and long-wavelength susceptibility. Adding a source term $\varphi(\vec{x}, t)$ to the right hand side of eq. (4) and going to the Fourier space one obtains the generalized response function

$$\tilde{G}(\vec{k}, \omega) = \left\langle \frac{\tilde{h}(\vec{k}, \omega)}{\tilde{\varphi}(\vec{k}, \omega)} \right\rangle_{\tilde{\varphi}=0} = \frac{1}{[\tilde{G}_0(\vec{k}, \omega)]^{-1} - \tilde{\Sigma}(\vec{k}, \omega)}, \quad (6)$$

where

$$[\tilde{G}_0(\vec{k}, \omega)]^{-1} = \Gamma \vec{k}^2 + i\lambda\omega + 4\mu_0 \mathcal{N} M_s^2 \hat{\delta}(\vec{k}, L) + \tilde{K}(\vec{k}). \quad (7)$$

is the bare correlator and $\tilde{\Sigma}(\vec{k}, \omega)$ is the "self-energy". $\hat{\delta}(\vec{k}, L)$ is the Fourier transform of L^{-d} . In the thermodynamic limit ($L \rightarrow \infty$) $\hat{\delta}(\vec{k}, L) \approx 1$ for $\vec{k} \rightarrow 0$ and zero otherwise. Since $\tilde{\Sigma}(0, 0) = 0$ and $\tilde{G}_0(0, 0)^{-1} = 4\mu_0 \mathcal{N} M_s^2$ it results that the low-frequency and long-wavelength susceptibility (or simply the susceptibility) is given by

$$\chi = \tilde{G}(0, 0) = \frac{1}{4\mu_0 \mathcal{N} M_s^2}. \quad (8)$$

This result is also exact to all orders of perturbation expansion.

In eq. (8) one cannot determine precisely which is the control parameter of the model. The susceptibility may diverges if both $\mathcal{N}$ or M_s goes to zero. To answer this question we look for a self-similar solution of eq. (4). Performing the scale transformation $x \rightarrow bx$, $L \rightarrow bL$, $t \rightarrow b^z t$ and $y \rightarrow b^\zeta y$ where z and ζ are the dynamic and roughness exponent, respectively, and taking into account eq. (5), eq. (4) becomes

$$\begin{aligned} \lambda b^{1-z} \frac{\partial}{\partial t} y(\vec{x}, t) &= \Gamma b^{-1} \nabla^2 y(\vec{x}, t) - \lambda b^{1-\zeta} v \\ &+ b^{1-\zeta} \eta[b\vec{x}, vb^z t + b^\zeta y(\vec{x}, t)] \\ &- 4\mu_0 \mathcal{N} M_s^2 b \int \frac{d^d x'}{L^d} y(\vec{x}', t) \\ &+ \int d^d x' K(\vec{x} - \vec{x}') [y(\vec{x}', t) - y\vec{x}, t)]. \end{aligned} \quad (9)$$

Notice that we cannot obtain scale invariance in the elastic and dipolar interaction terms simultaneously. Thus we have to determine which of this terms gives the major contribution. A magnitude which characterizes the ration between elastic and dipolar interactions is

$$r = \frac{\mu_0 M_s^2}{4\pi^2 \Gamma \Lambda}, \quad (10)$$

where Λ , the momentum cutoff, is of the order the inverse of the domain wall thickness. The case $r \ll 1$ where dipolar interactions can be neglected has been already analyzed in [17], here we focus our attention in the opposite case $r \gg 1$ where dipolar interactions are relevant. In this case, imposing scale invariance we obtain that $\mathcal{N}$ and v should go to zero and

$$z = 1, \quad \zeta = \frac{2-d}{2}. \quad (11)$$

The scaling exponents obtained in this way are identical to those derived by CZDE [10] for the case of constant magnetic field. Moreover the upper critical dimension $d_c = 2$ is also the same.

The saturation magnetization is not a control parameter because the dipolar interaction term, which is proportional to M_s^2 , does not renormalize. Thus, $\mathcal{N}$ and v are the only control parameters. Since scale invariance is obtained when $\mathcal{N} = 0^+$ from eq. (9) one can define the correlation length $\xi \sim \mathcal{N}^{-\nu}$ with $\nu = 1$. Moreover, from eq. (8) we obtain that the susceptibility scale as $\chi \sim \mathcal{N}^{-\gamma}$ with $\gamma = 1$. At the critical state $\xi \sim L$ and

$$\chi \sim \xi^{\gamma\nu} \sim L, \quad (12)$$

On the other hand, the increase of v may change the character of the noise correlator. For large v the noise correlator can be approximated by an annealed noise. In this case the magnetic field predominates over disorder, corresponding with a supercritical regime. From eq. (9) one can define the characteristic velocity $v_c \sim \epsilon^\theta$ with

$$\theta = \nu(z - \zeta), \quad (13)$$

which divides the phase diagram $(\mathcal{N}, v)$ into two regions, the supercritical state $v \gg v_c$ and the subcritical one $v \ll v_c$. Criticality is obtained when $\mathcal{N} \rightarrow 0$ and $v \rightarrow 0$.

To go further we have performed a RG analysis of the problem. We integrate out the degrees of freedom in a momentum shell near the cutoff Λ and rescale $k \rightarrow b^{-1}k$, $\omega \rightarrow b^{-z}\omega$, and $\tilde{w} \rightarrow b^{\zeta+d+z}\tilde{w}$, where $b = e^l$ with $l \rightarrow 0$. The flow equations for the parameters Γ , λ , $\mathcal{N}$, M_s and v are obtained through a direct application of the RG transformations to eq. (4) in the Fourier space. The renormalization of the moments of the noise correlator ($Q_n = \int_q \tilde{\Delta}(q) q^n$) is obtained considering vertex functions [16,15]. As in perturbation theory here we only consider the case $r \gg 1$. In this case we found corrections of the order of r^{-1} to the exponents computed by perturbation theory [18]. Thus, the exponents z and ζ in eq. (11) are exact up to $\mathcal{O}(r^{-1})$.

Let us now determine the avalanche distribution exponents. The Barkhausen signal $V(t)$ is the voltage produced from a pickup coil around a ferromagnet subjected

to a slowly varying applied field. In the low-frequency limit the time scale for domain wall motion is much smaller than the time between jumps and, therefore, one may guarantee that each induced voltage jump corresponds with a single avalanche in the domain wall motion. A resolution voltage level V_R is defined, such that one can not resolve details below V_R . An elementary Barkhausen jump can thus be defined as the portion of the $V(t)$ signal delimited by two subsequent intersections of the signal with the V_R line. With this definition, the duration T is simply the time interval between these two subsequent intersections and the size s is the area delimited by $V(t)$ and V_R between the same points.

In the subcritical regime the dynamics takes place in the form of avalanches, characterized by the avalanche size $P(s) = s^{-\tau} f(s/s_c)$ and duration $P(T) = T^{-\alpha} g(T/T_c)$ distributions, where s_c and T_c are the avalanche size and duration cutoffs. In the subcritical state $s_c \sim \epsilon^{-1/\sigma}$ and $\xi \sim \epsilon^{-\nu}$ while at criticality $s_c \sim L^D$ and $\xi \sim L$, where D is the avalanche dimension, leading to the scaling relation

$$\sigma = \frac{1}{D\nu}. \quad (14)$$

Other scaling relations are obtained taking into account that $\chi = \langle s \rangle$ and $\int ds P(s) = \int dT P(T)$, which lead to

$$\gamma = \frac{2-\tau}{\sigma}, \quad (\tau-1)D = (\alpha-1)z, \quad (15)$$

respectively.

Using these scaling relations and our result $\gamma = \nu = z = 1$ we obtain the following scaling relations for the avalanche exponents

$$\tau = 2 - \frac{1}{\alpha}, \quad \alpha = D, \quad (16)$$

On the other hand, for $d < d_c$, the avalanche dimension and the roughness exponent are related via $D = d + \zeta$, while above the upper critical dimension one obtains $D = d_c = 2$. Hence, we can compute τ and α using eq. (16) and this value of D . For instance

$$\begin{aligned} \alpha &= \frac{3}{2}, \text{ for } d+1=2; \\ &= 2, \text{ for } d+1 \geq 3. \end{aligned} \quad (17)$$

The scaling law in eq. (12), the scaling relations in eq. (16) and the values of the scaling exponent α in eq. (17) are also obtained for DASM [14]. However, the upper critical dimension of DASM is 3, and not 2 as obtained here. This apparent contradiction can be understood if one takes into account that in DASM time evolution does not introduce an additional dimension because it can be represented by the evolution in the preferential dimension. On the contrary in the motion of the domain wall the time evolution take place perpendicular to the d -dimensional substrate and, therefore, introduces

an additional dimension. Taking this fact into account the upper critical dimension will be $2 + 1$, as in DASM. Thus, the CZDE model in $d + 1$ dimensions is equivalent to a d -dimensional DASM.

The fact that the CZDE is mapped into a directed sandpile model, and not into an undirected one, is due to dipolar interactions which introduces an anisotropy in the system. If dipolar interactions becomes negligible ($r \ll 1$) then we expect that the CZDE model will be mapped into an undirected sandpile model. Experiments in magnetostrictive materials, where dipolar interactions can be neglected, have been performed [12,19,20]. Earlier measurements by Urbach *et al* [12] gives $\tau = 1.33 \pm 0.10$. More recently, Durin and Zapperi (DZ) [20] reported the more accurate exponents $\tau = 1.28 \pm 0.02$ and $\alpha = 1.5 \pm 0.1$. On the other hand, numerical simulations of the CZDE model in two dimensions and without dipolar interactions have also been performed [19,20]. The more accurate numerical estimates, reported by Durin and Zapperi [20], are $\tau = 1.26 \pm 0.04$ and $\alpha = 1.40 \pm 0.05$. These exponents are in the range reported for undirected Abelian sandpile models in two dimensions. For instance, numerical simulations of the Bak-Tang-Wiesenfeld (the prototype of undirected Abelian sandpile model) gives $\tau = 1.293$ and $\alpha = 1.480$ [21]. However, to validate this guess we cannot base our analysis only in the avalanche exponents, we must provide further elements. In [17] we have shown that the CZDE without dipolar interactions can be mapped into a class of undirected sandpile models with annealed noise in the toppling rule.

In conclusion we have shown that the CZDE model for the dynamics of a domain wall belongs to the universality class of directed Abelian sandpile models. Dipolar interactions introduce an anisotropy in the systems leading to a preferential direction along the magnetization. We have provided strong arguments which states the equivalence between the dynamics of a domain wall and sandpile models. This conclusion can be extended to systems with many domain walls, since for short time scales the interaction between domain walls can be neglected and the single domain wall picture is correct. Hence, the Barkhausen effect actually exhibits self-organized critical behavior.

-
- [1] P. Bak, C. Tang, and K. Wiesenfeld, Phys. Rev. Lett. **59**, 381 (1987); Phys. Rev. A **38**, 364 (1988).
 - [2] A. Vespignani and S. Zapperi, Phys. Rev. Lett. **78**, 4793 (1997); Phys. Rev. E **57**, 6345 (1998).
 - [3] O. Geoffroy and J. L. Corteseil, J. Magn. Magn. Mate. **97**, 198 (1991); *ibid* **97**, 205 (1991).
 - [4] P. J. Cote and L. V. Meisel, Phys. Rev. Lett. **67**, 1334 (1991).
 - [5] K. P. O'Brien and M. B. Weissman, Phys. Rev. E **50**, 3446 (1994).
 - [6] D. Spasojević, S. Buknić, S. Milošević, and H. E. Stanley, Phys. Rev. E **54**, 2531 (1996).
 - [7] J. P. Sethna, K. Dahmen, S. Kartha, J. Krumhansl, B. W. Roberts, and J. D. Shore, Phys. Rev. Lett. **21**, 3347 (1993).
 - [8] O. Percović, K. Dahmen, and J. P. Sethna, Phys. Rev. Lett. **75**, 4528 (1995).
 - [9] K. Dahmen and J. Sethna, Phys. Rev. Lett. **71**, 3222 (1993); Phys. Rev. B **53**, 14872 (1996).
 - [10] P. Cizeau, S. Zapperi, G. Durin, and H. E. Stanley, Phys. Rev. Lett. **79**, 4669 (1997); S. Zapperi, P. Cizeau, G. Durin, and H. E. Stanley, Phys. Rev. B, 1998.
 - [11] B. Alessandro, C. Beatrice, G. bertotti, and A. Montorsi, J. Appl. Phys. **68**, 2901 (1990); *ibid* **68**, 2908 (1990).
 - [12] J. S. Urbach, R. C. Madison, and J. T. Markert, Phys. Rev. Lett. **75**, 276 (1995).
 - [13] When dipolar interactions are negligible a different universality class is obtained.
 - [14] For a review see D. Dhar, cond-mat/9808047.
 - [15] O. Narayan and D. S. Fisher, Phys. Rev. Lett. **68**, 3615 (1992); Phys. Rev. B **48**, 7030 (1993).
 - [16] T. Nattermann, S. Stepanow, L.-H. Tang, and H. Leschhorn, J. Phys. II France **2**, 1483 (1992); H. Leschhorn, T. Nattermann, S. Stepanow, and L.-H. Tang, Ann. Phys. **6**, 1 (1997).
 - [17] A. Vázquez and O. Sotolongo-Costa, cond-mat/9812173.
 - [18] A. Vázquez and O. Sotolongo-Costa, in progress.
 - [19] M. Bahiana, B. Koiller, S. L. A. de Queiroz, J. C. Denardin, and R. L. Sommer, cond-mat/9808017.
 - [20] G. Durin and S. Zapperi, cond-mat/9808224.
 - [21] S. Lübeck and K. Usadel, Phys. Rev. E **55**, 4095 (1997).